

Charm Antiquark and Charm Quark in the Nucleon*

Xiaotong Song*

*Institute of Nuclear and Particle Physics
Jesse W. Beams Laboratory of Physics
Department of Physics, University of Virginia
Charlottesville, VA 22904, USA*

We estimate the intrinsic charm contributions to the quark flavor and spin observables of the nucleon in the SU(4) quark meson fluctuation model. In this model, the charm or anticharm reside in the charmed mesons created by the nonperturbative quantum quark-meson fluctuations. The intrinsic charm content in the proton, $2\bar{c}/\sum(q+\bar{q}) \simeq 0.011 \pm 0.008$, is almost one order of magnitude smaller than the intrinsic strange content. The intrinsic charm helicity is also small and negative, $\Delta c \simeq -(0.009 \pm 0.006)$. The fraction of the total quark helicity carried by the charm is $|\Delta c/\Delta\Sigma| \simeq 0.021 \pm 0.014$. The ratio of the charm with positive helicity to that with negative helicity is $c_{\uparrow}/c_{\downarrow} = 35/67$. For the intrinsic strange component, one has $s_{\uparrow}/s_{\downarrow} \simeq 7/13$. A detail comparison of our predictions with data and other models or analyses is given. The intrinsic charm contribution to the Ellis-Jaffe sum rule is also discussed.

PACS numbers: 14.65.Dw, 14.20.Dh, 12.39.Fe, 11.30.Hv

1. Introduction

As suggested by many authors long time ago [1, 2], there are so called ‘*intrinsic*’ heavy quark components in the proton wave function. The intrinsic heavy quarks are created from the quantum fluctuations associated with the bound state hadron dynamics. They exist in the hadron over a long time independent of any external probe momentum transferred in the collision. The probability of finding the intrinsic heavy quarks in the hadron is completely determined by nonperturbative mechanisms. On the other hand, the *extrinsic* heavy quarks are created on a short time scale in association with a large transverse momentum reaction and their distributions can be derived from QCD bremsstrahlung and pair production processes, which lead to standard QCD evolution. The study of nonperturbative mechanism $gg \rightarrow q\bar{q}$ [1] shows that the intrinsic heavy quark contribution scales as $1/m_Q^2$, where m_Q is the mass of the heavy quark, the probability of finding the intrinsic bottom is expected to be much smaller, hence we will only consider the intrinsic charm (IC) component, e.g. $|uudc\bar{c}\rangle$, in the nucleon.

Many works on intrinsic charm are based on models of the nucleon (see e.g. MIT bag model [2], NJL model [3], meson cloud model [1, 4], instanton model [5, 6] and others [7], etc.) or combination of the IC hypothesis and analysing the existing DIS data on charm production to obtain some information of the intrinsic charm content (see e.g. [8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18]). As pointed out in [15] that a non-perturbative IC component was unable to explain enhancement in the e^+p neutral current cross section at HERA kinematics, but HERA

data may provide a good test of the intrinsic charm if the charm component of F_2 is directly isolated and no new physics exists. Another discussion given in [16] suggested a nonsymmetric charm distribution, for which $\bar{c}$ is much harder than c , which may account for the HERA anomaly, a significant enhancement of cross sections at HERA at large x and Q^2 [19, 20]. However, the most recent analysis given in [17], which includes both the perturbative photon-gluon fusion process and nonperturbative intrinsic charm contribution, found no conclusive evidence for the intrinsic charm and only provided an upper bound of around 0.4%. Therefore what is the effect of charm in the structure of the proton is still an open and interesting question. A comprehensive review on the nucleon sea, which includes light quark, intrinsic strange and charm, can be found in [21].

The chiral quark model or more precisely the quark meson fluctuation model (in SU(2) version) was first suggested in [22] to explain the sea flavor asymmetry, $\bar{d} - \bar{u} > 0$. The model was extended to SU(3) without and with SU(3) symmetry breaking to describe the quark spin structure (see e.g. [23, 24, 25, 26, 27] and references therein), and orbital structure (see e.g. [28] and references therein) of the nucleon. In this model, the nucleon structure is determined by its zeroth order valence quark configuration and all possible quantum fluctuations of the valence quark into a recoil quark and a meson. It provides a natural and nonperturbative explanation of a nonzero (intrinsic) strange sea and negative strange sea polarization through the quark-kaon fluctuation (e.g. $u \rightarrow s + K^+$). The results agree quite well with the data obtained in deep-inelastic scattering and other related experiments.

Based on earlier encouraging result [27], we suggested an SU(4) version of quark-meson fluctuation model and predicted a nonzero intrinsic charm in the proton [29]. In the SU(4) version, charm or anti-charm quarks re-

*Electronic address: xs3e@virginia.edu;

* Report No. INPP-UVA-2001-04 (September 18, 2001)

side in the charmed mesons which are created by non-perturbative quantum quark-meson fluctuations such as $u \rightarrow c + \bar{D}^0$. These charm or anti-charm quarks are essentially *intrinsic* components of the hadron.

The main motivations of extending to SU(4) version are: 1) There is no physical reason to forbid the fluctuation from a light quark to a charmed quark and charmed meson. Hence it is interesting to investigate what are the changes of the SU(3) predictions after including the charm or anticharm contributions. 2) According to the symmetric GIM model [30], one should deal with the weak axial current in the framework of SU(4) symmetry. It implies that the charm quark should play some role in determining the spin, flavor and orbital structure of the nucleon. Therefore it is natural and reasonable to extend the SU(3) version to SU(4) by including the charm and anticharm contributions.

In this paper, we will discuss the effect of the IC contribution to the flavor and spin observables of the proton in the SU(4) quark meson fluctuation model with symmetry-breaking. The analytic and numerical results of the flavor and spin contents for each quark flavor u , d , s , and c in the proton are presented. Predictions are compared with the existing data and other model results or analyses. The intrinsic charm contribution to the Ellis-Jaffe sum rule is also discussed.

2. SU(4) model with symmetry breaking

In the framework of SU(4) quark model, there are sixteen pseudoscalar mesons, a 15-plet and a singlet. In this paper, the contribution of the SU(4) singlet will be neglected. It is easy to show that the effective Lagrangian describing interaction between quarks and the mesons in the SU(4) case is

$$L_I = g_{15} \bar{q} \begin{pmatrix} G_u^0 & \pi^+ & \sqrt{\epsilon} K^+ & \sqrt{\epsilon_c} \bar{D}^0 \\ \pi^- & G_d^0 & \sqrt{\epsilon} K^0 & \sqrt{\epsilon_c} D^- \\ \sqrt{\epsilon} K^- & \sqrt{\epsilon} \bar{K}^0 & G_s^0 & \sqrt{\epsilon_c} D_s^- \\ \sqrt{\epsilon_c} D^0 & \sqrt{\epsilon_c} D^+ & \sqrt{\epsilon_c} D_s^+ & G_c^0 \end{pmatrix} q, \quad (1)$$

where $D^+ = (c\bar{d})$, $D^- = (\bar{c}d)$, $D^0 = (c\bar{u})$, $\bar{D}^0 = (\bar{c}u)$, $D_s^+ = (c\bar{s})$, and $D_s^- = (\bar{c}s)$. The neutral charge components $G_{u(d)}^0$ and $G_{s,c}^0$ are defined as

$$G_{u(d)}^0 = +(-)\pi^0/\sqrt{2} + \eta^0 \sqrt{\epsilon_\eta/6} + \eta'^0 \sqrt{\zeta'^2/48} - \eta_c^0 \sqrt{\epsilon_c/16} \quad (2)$$

$$G_s^0 = -\eta^0 \sqrt{2\epsilon_\eta/3} + \eta'^0 \sqrt{\zeta'^2/48} - \eta_c^0 \sqrt{\epsilon_c/16} \quad (3)$$

$$G_c^0 = -\eta'^0 \sqrt{3\zeta'^2/16} + \eta_c^0 \sqrt{9\epsilon_c/16} \quad (4)$$

with

$$\pi^0 = (u\bar{u} - d\bar{d})/\sqrt{2}, \quad \eta^0 = (u\bar{u} + d\bar{d} - 2s\bar{s})\sqrt{6}, \quad (5)$$

$$\eta'^0 = (u\bar{u} + d\bar{d} + s\bar{s})/\sqrt{3}, \quad \eta_c^0 = (c\bar{c}). \quad (6)$$

Similar to the SU(3) case, we define $a \equiv |g_{15}|^2$, which denotes the transition probability of splitting $u(d) \rightarrow d(u) +$

$\pi^{+(-)}$. Hence ϵa , $\epsilon_\eta a$, $\zeta'^2 a$, and $\epsilon_c a$ denote the probabilities of splittings $u(d) \rightarrow s + K^{-(0)}$, $u(d) \rightarrow u(d) + \eta^0$, $u(d) \rightarrow u(d) + \eta'^0$ and $u(d) \rightarrow c + \bar{D}^0 (D^-)$ respectively. If the splitting probability is dominated by the mass effects, we expect $0 < \epsilon_c a < \zeta'^2 a < \epsilon a \simeq \epsilon_\eta a < a$, or

$$0 < \epsilon_c < \zeta'^2 < \epsilon \simeq \epsilon_\eta < 1. \quad (7)$$

It implies that the probability of emitting heavier meson such as D from light quarks is smaller than that of emitting the lighter mesons such as K , η , and η' , etc. Since the charm contributions only appeared in the observables which are explicitly related to the charm or anticharm variables, therefore we expect that the SU(4) model will not significantly change the SU(3) predictions on observables which are not explicitly charm-related. On the other hand, the SU(4) version leads to many *new* predictions on observables which are directly related to the charm or anticharm. These predictions can be tested by the future experiments. Before going to further discussion, we would like to make a few remarks.

- The quark contents of the η^0 and η'^0 shown in Eqs.(5) and (6) imply that they are actually η_8 and η_1 . Hence the octet-singlet mixing is neglected in our description.
- We have shown in [27] that a better SU(3) description can be achieved by taking $\epsilon_\eta \simeq \epsilon$ and $\zeta'^2 < 1$, i.e. the effective U(1)-breaking parameter is smaller than other SU(3)-breaking parameters. This result was deduced from the data of $\bar{d} - \bar{u}$, which requires ζ' should be small and negative. More detail discussion can be found in section II and Fig.1 in [27]. It can be shown that including the charm contributions will not change this conclusion. Therefore, we may choose $\zeta'^2 = 0$ as one possible option in the numerical calculation, i.e. neglect the singlet contribution from the beginning. In this case, the number of independent parameters would be only *three*: a , ϵ , and ϵ_c . Since our analytical formulae are not restricted to this approximation, we can also discuss $\zeta'^2 \neq 0$ case (see discussion in section 4 below).
- It should be noted that the definition of $G_{u(d)}^0$ in SU(4) case is different from that in the SU(3) case. Besides an additional *charm* term, $-\eta_c^0 \sqrt{\epsilon_c/16}$, the coefficient of η^0 term is changed. For G_s^0 , even the coefficient of η'^0 term is also changed. Therefore, the SU(4) formalism cannot be reduced to SU(3) simply by taking $\epsilon_c \rightarrow 0$ only [see later discussion on Eqs. (15) and (16)]. The G_c^0 is completely new in the SU(4) version.
- As pointed out in the original chiral quark model [22, 23, 24, 25, 26, 27], the nucleon properties are defined in the scale range between Λ_{QCD} ($\sim 0.2 - 0.3$ GeV) and Λ_{XSB} (~ 1 GeV), where the spontaneous breaking of chiral symmetry leads to

the existence of Goldstone bosons. In the quark meson fluctuation model (SU(3) in [28] and SU(4) in this paper), we will assume all calculated quantities are also defined in the same scale range $[(0.2 \text{ GeV})^2 < \mu^2 < (1.0 \text{ GeV})^2]$. At this μ^2 range, the sea content should be dominated by the intrinsic component created by nonperturbative mechanism, quark-meson fluctuation, [31] and the extrinsic sea component is expected to be small.

- We note that in our formalism, only the integrated flavor content $[q(Q^2) = \int_0^1 dx q(x, Q^2)]$ and helicity content $[\Delta q(Q^2) = \int_0^1 dx \Delta q(x, Q^2)]$ are discussed. To make brief of the formalism, we will omit the argument Q^2 everywhere unless specified otherwise.

In addition to the allowed fluctuations discussed in the SU(3) case, a valence quark is now allowed to split up or fluctuate to a recoil charm quark and a charmed meson. For example, a valence u-quark with spin-up, the allowed fluctuations are

$$u_\uparrow \rightarrow d_\downarrow + \pi^+, \quad u_\uparrow \rightarrow s_\downarrow + K^+, \quad u_\uparrow \rightarrow u_\downarrow + G_u^0, \quad (8)$$

$$u_\uparrow \rightarrow c_\downarrow + \bar{D}^0, \quad (9)$$

$$u_\uparrow \rightarrow u_\uparrow. \quad (10)$$

Similarly, one can list the allowed fluctuations for $u_\downarrow$, $d_\uparrow$, $d_\downarrow$, $s_\uparrow$, and $s_\downarrow$. Since we are only interested in the spin-flavor structure of the nucleon (or other non-charmed hadrons), which does not have valence charm quark, hence the fluctuations from a valence charm quark will not be discussed.

The spin-up and spin-down quark or antiquark contents in the proton, up to first order of the quantum fluctuation, can be written as (a detail SU(3) version see e.g. [28]),

$$n_p(q'_{\uparrow,\downarrow}, \text{ or } \bar{q}'_{\uparrow,\downarrow}) = \sum_{q=u,d,s,c} \sum_{h=\uparrow,\downarrow} n_p^{(0)}(q_h) P_{q_h}(q'_{\uparrow,\downarrow}, \text{ or } \bar{q}'_{\uparrow,\downarrow}), \quad (11)$$

where $q' = u, d, s, c$ and $n_p^{(0)}(q_{\uparrow,\downarrow})$ are determined by the zeroth order, i.e. naive quark model (NQM), valence quark wave function of the proton and

$$n_p^{(0)}(u_\uparrow) = 5/3, \quad n_p^{(0)}(u_\downarrow) = 1/3, \quad (12)$$

$$n_p^{(0)}(d_\uparrow) = 1/3, \quad n_p^{(0)}(d_\downarrow) = 2/3.$$

In Eq.(11), $P_{q_{\uparrow,\downarrow}}(q'_{\uparrow,\downarrow})$ and $P_{q_{\uparrow,\downarrow}}(\bar{q}'_{\uparrow,\downarrow})$ are probabilities of finding a quark $q'_{\uparrow,\downarrow}$ or an antiquark $\bar{q}'_{\uparrow,\downarrow}$ arise from all quantum fluctuations of a valence quark $q_{\uparrow,\downarrow}$. The probabilities $P_{q_{\uparrow,\downarrow}}(q'_{\uparrow,\downarrow})$ and $P_{q_{\uparrow,\downarrow}}(\bar{q}'_{\uparrow,\downarrow})$ can be obtained from the effective Lagrangian (1) and have been listed in Table I, where only $P_{q_\uparrow}(q'_{\uparrow,\downarrow})$ and $P_{q_\uparrow}(\bar{q}'_{\uparrow,\downarrow})$ are shown. Those arise from $q_\downarrow$ can be obtained by using $P_{q_\downarrow}(q'_{\uparrow,\downarrow}) =$

$P_{q_\uparrow}(q'_{\downarrow,\uparrow})$ and $P_{q_\downarrow}(\bar{q}'_{\uparrow,\downarrow}) = P_{q_\uparrow}(\bar{q}'_{\downarrow,\uparrow})$. The notations appeared in Table I are defined as

$$f \equiv 1/2 + \epsilon_\eta/6 + \zeta'^2/48 + \epsilon_c/16, \quad (13)$$

$$f_s \equiv 2\epsilon_\eta/3 + \zeta'^2/48 + \epsilon_c/16,$$

and

$$\tilde{A} \equiv 1/2 - \sqrt{\epsilon_\eta}/6 - \zeta'/12, \quad \tilde{B} \equiv -\sqrt{\epsilon_\eta}/3 + \zeta'/12, \quad (14)$$

$$\tilde{C} \equiv 2\sqrt{\epsilon_\eta}/3 + \zeta'/12, \quad \tilde{D} \equiv \sqrt{\epsilon_c}/4.$$

Analogous to the SU(3) case, the special combinations $\tilde{A}$, $\tilde{B}$, $\tilde{C}$, and $\tilde{D}$ stem from the definitions (2)-(4), which show the quark and antiquark contents in the neutral bosons $G_{u,d,s,c}^0$. The numbers f and f_s stand for the probabilities of the quark splitting $u_\uparrow(d_\uparrow) \rightarrow u_\downarrow(d_\downarrow) + G_{u(d)}^0$ and $s_\uparrow \rightarrow s_\downarrow + G_s^0$ respectively.

To reduce to the SU(3) version, one should take the limit $\epsilon_c \rightarrow 0$ and change ζ' to $4\zeta'$ simultaneously. This is because we have different neutral combinations $G_{u,d,s}^0$ from the SU(3) ones. This difference shows that the *charm* contribution is not only presented in the *charm* term which is proportional to ϵ_c , but also affected by the ζ' term. Taking $\epsilon_c \rightarrow 0$ and $\zeta' \rightarrow 4\zeta'$, one can see from Eq. (13) that the f and f_s indeed reduce to the corresponding quantities in the SU(3) case,

$$(f)_{SU(3)} = 1/2 + \epsilon_\eta/6 + \zeta'^2/3, \quad (15)$$

$$(f_s)_{SU(3)} = 2\epsilon_\eta/3 + \zeta'^2/3,$$

which is exactly Eq.(6b) in [28]. Taking $\epsilon_c \rightarrow 0$ and $\zeta' \rightarrow 4\zeta'$, one has $\tilde{D} \rightarrow 0$ and

$$\tilde{A} \rightarrow A_{SU(3)}/3, \quad \tilde{B} \rightarrow B_{SU(3)}/3, \quad \tilde{C} \rightarrow C_{SU(3)}/3, \quad (16)$$

where $A_{SU(3)}$, $B_{SU(3)}$, and $C_{SU(3)}$ are the same as A , B , and C in Eq. (6a) in [28].

If Eq.(12) is replaced by $n_n^{(0)}(d_\uparrow) = 5/3$, $n_n^{(0)}(d_\downarrow) = 1/3$, $n_n^{(0)}(u_\uparrow) = 1/3$, and $n_n^{(0)}(u_\downarrow) = 2/3$, from (11), we will obtain all flavor and spin contents in the neutron.

3. Flavor and spin contents in the proton

We note that the quark *flips* its spin in the quark splitting processes $q_{\uparrow,\downarrow} \rightarrow q_{\downarrow,\uparrow} + \text{meson}$, i.e. processes in (8) and (9), but not in the (10) (*no splitting*). The quark helicity non-flip contribution in splitting processes is entirely neglected. This is a basic assumption in the model and seems to be consistent with the picture given by the instanton model.

3.1. Quark flavor content

Using (11), (12) and the probabilities $P_{q_{\uparrow,\downarrow}}(q'_{\uparrow,\downarrow})$ and $P_{q_{\uparrow,\downarrow}}(\bar{q}'_{\uparrow,\downarrow})$ listed in Table I, one obtains all quark and antiquark flavor contents in the proton,

$$u = 2 + \bar{u}, \quad d = 1 + \bar{d}, \quad s = 0 + \bar{s}, \quad c = 0 + \bar{c}, \quad (17)$$

where

$$\bar{u} = a[1 + \tilde{A}^2 + 2(1 - \tilde{A})^2], \quad \bar{d} = a[2(1 + \tilde{A}^2) + (1 - \tilde{A})^2], \quad (18)$$

$$\bar{s} = 3a[\epsilon + \tilde{B}^2], \quad \bar{c} = 3a[\epsilon_c + \tilde{D}^2]. \quad (19)$$

From (18), one obtains

$$\bar{u}/\bar{d} = 1 - 6\tilde{A}/[(3\tilde{A} - 1)^2 + 8], \quad (20)$$

and

$$\bar{d} - \bar{u} = 2a\tilde{A}. \quad (21)$$

Similarly, one can obtain $2\bar{c}/(\bar{u} + \bar{d})$, $2\bar{c}/(u + d)$, $2\bar{c}/\sum(q + \bar{q})$ and other flavor observables related to charm quark and charm antiquark. We note that in the SU(4) case, one has

$$\sum_{q, \bar{q}} (q + \bar{q}) = 3(1 + 2a\xi_1), \quad (22)$$

where $\xi_1 \equiv 1 + \epsilon + \epsilon_c + f$. It is easy to verify that in the limit $\epsilon_c \rightarrow 0$, and using Eq.(16), all SU(4) results reduce to those given in the SU(3) case. Two remarks on (20) and (21) are in order.

- Defining the ratio, $r \equiv \bar{u}/\bar{d}$, we obtain from Eq. (20) that $\tilde{A} = 2 - r \pm \sqrt{14r - 8r^2 - 5}$. Since $\tilde{A}$ must be a *real* number due to (21), hence $14r - 8r^2 - 5$ must be positive or zero, and the ratio r satisfies

$$1/2 \leq \bar{u}/\bar{d} \leq 5/4, \quad (23)$$

which seems to be consistent with the experimental data shown in Table III.

- Since $|\zeta'/2| \leq \sqrt{\epsilon_\eta}$ in our approximation, from (14), we have $\tilde{A} \leq 1/2$ (note that ζ' is *negative*) and

$$\bar{d} - \bar{u} \leq a. \quad (24)$$

It gives a lower bound of the parameter a – the probability of quark splitting $q \rightarrow q' + \pi$.

3.2. Quark helicity content

Similar to the SU(3) version, we have

$$\Delta q = \sum_{q'=u,d} [n^{(0)}(q'_\uparrow) - n^{(0)}(q'_\downarrow)][P_{q'_\uparrow}(q_\uparrow) - P_{q'_\uparrow}(q_\downarrow)]. \quad (25)$$

Using Eqs. (12), (25), and Table I, we obtain

$$\Delta u = 4[1 - a(\epsilon + \epsilon_c + 2f)]/3 - a, \quad (26)$$

$$\Delta d = -[1 - a(\epsilon + \epsilon_c + 2f)]/3 - a, \quad (27)$$

$$\Delta s = -a\epsilon, \quad (28)$$

$$\Delta c = -a\epsilon_c, \quad (29)$$

$$\Delta \Sigma \equiv \sum_{q=u,d,s,c} \Delta q = 1 - 2a(1 + \epsilon + \epsilon_c + f), \quad (30)$$

or

$$\Delta \Sigma/2 = 1/2 - a\xi_1, \quad (31)$$

where ξ_1 is defined in (22), and

$$\Delta \bar{q} = 0, \quad (\bar{q} = \bar{u}, \bar{d}, \bar{s}, \bar{c}). \quad (32)$$

Before going to the numerical calculation, we would like to make several remarks on some results which are basically independent of parameters.

- Comparing with the SU(3) case, a new ϵ_c term, which presents the intrinsic charm contribution, appeared in Δu , Δd , Δc , and $\Delta \Sigma$, but not in Δs . This is because there is no process which can mix the *strange* helicity and *charm* helicity contributions. Δs can only come from the processes like $u_\uparrow \rightarrow s_\downarrow + K^+$, while Δc comes only from the processes like $u_\uparrow \rightarrow c_\downarrow + \bar{D}^0$. Although there are s , $\bar{s}$ (or c , $\bar{c}$) in all *neutral* bosons $G_{u,d,s,c}^0$, they give *no* contributions to Δs (or Δc) due to $s_\uparrow$ and $s_\downarrow$ (or $c_\uparrow$ and $c_\downarrow$) appeared with *equal probability* in these neutral bosons.
- The charm quark helicity Δc , (29), is *nonzero* as far as ϵ_c is nonzero. Analogous to the strange quark helicity, Δc is definitely *negative*, but the size of the intrinsic charm helicity depends on the parameter ϵ_c and the splitting probability a .
- The physical meaning of (31) is that the total *loss* of the quark helicity arises from *four* splitting processes with quark spin-flip, *three* in (8) and *a new* splitting in (9). Comparing with the SU(3) case, where $\Delta \Sigma/2 = 1/2 - a(1 + \epsilon + f)$, we now have an additional reduction, $-a\epsilon_c$, of the total quark spin due to the splitting related to the charm.
- In the splitting process $u_{\uparrow(\downarrow)} \rightarrow c_{\downarrow(\uparrow)} + \bar{D}^0$, the anticharm resides only in the charmed meson, e.g. $\bar{D}^0(\bar{c}, u)$. The probabilities of finding $\bar{c}_\uparrow$ and $\bar{c}_\downarrow$ are equal in the spinless charmed meson. Therefore $\Delta \bar{c} = 0$. Similar discussion in the SU(3) case [24] has led to $\Delta \bar{q} = 0$ for $\bar{q} = \bar{u}, \bar{d}, \bar{s}$. The result (32) shows that the helicities of the sea quark and antiquark are not equal, $\Delta q_{sea} \neq \Delta \bar{q}$. This is different from the usual gluon splitting ($g \rightarrow q + \bar{q}$) model and $gg \rightarrow q + \bar{q}$ model [1]. In the gluon splitting and gluon fusion models, the sea quark and antiquark with the same flavor are perturbatively or nonperturbatively created as a *pair* from the gluon or gluons and $\Delta q_{sea} = \Delta \bar{q}$. The DIS data [42] seems to support the prediction $\Delta \bar{q} \simeq 0$ but with large errors.

- From (19) and (29), using $\tilde{D}^2 = \epsilon_c/16$ in (14), one can see that the ratio

$$\Delta c/c = -16/51 \quad (33)$$

is a constant *independent* of any splitting parameters. This is a special prediction for the charm flavor in the SU(4) quark meson model.

- For the strange flavor, from $\tilde{B}$ in (14) and $\bar{s}$ in (19), one has $s = \bar{s} \simeq \epsilon a(10/3)[1 + |\zeta'|/(20\sqrt{\epsilon})]$ due to $\zeta'^2 \ll 1$. In the limit $\zeta'^2 \rightarrow 0$, we have

$$\Delta s/s = -3/10. \quad (34)$$

Hence $\Delta s/s$ is also a constant in this limit.

- From (33) and (32), one obtains the ratio of the charm with positive helicity to that with negative helicity is also a constant,

$$c_\uparrow/c_\downarrow = 35/67 \simeq 0.522. \quad (35)$$

It shows that more $c_\downarrow$ is created than $c_\uparrow$ in the splitting processes $u_{\uparrow(\downarrow)} \rightarrow c_{\downarrow(\uparrow)} + \bar{D}^0$ and $d_{\uparrow(\downarrow)} \rightarrow c_{\downarrow(\uparrow)} + D^-$. This is because the total contribution from $u_\uparrow$ and $d_\uparrow$ splittings is larger than that from $u_\downarrow$ and $d_\downarrow$ splittings. Similar situation occurs for the strange quarks. From (34) and (32), we have, for $\zeta' = 0$,

$$s_\uparrow/s_\downarrow = 7/13 \simeq 0.538. \quad (36)$$

- For the u -flavor and d -flavor, we do not have similar exact relations like (33) and (34). This is because the quantities $u_\uparrow - u_\downarrow$ and $d_\uparrow - d_\downarrow$ depend on ϵ_c , while $u_\uparrow + u_\downarrow$ and $d_\uparrow + d_\downarrow$ do not. Hence $\Delta u/u$ and $\Delta d/d$ depend on all three splitting parameters. The numerical calculation (see next section) shows that

$$\Delta u/u \simeq 0.383, \quad \Delta d/d \simeq -0.287. \quad (37)$$

From (37) and (32), we obtain

$$d_\uparrow/d_\downarrow \simeq 0.554, \quad (38)$$

and

$$u_\uparrow/u_\downarrow \simeq 2.241. \quad (39)$$

Comparing with the zeroth approximation (NQM), where $d_\uparrow/d_\downarrow = 1/2$ and $u_\uparrow/u_\downarrow = 5/1$, hence the quark-meson splittings lead to a small enhancement for $d_\uparrow/d_\downarrow$, but a significant reduction for $u_\uparrow/u_\downarrow$.

- The ratios of the flavor or spin contents predicted in this paper are also defined at $Q^2 = \mu^2$ scale, where the sea (light or heavy) quark contents are ‘intrinsic’ and determined by nonperturbative mechanism. To compare the predictions with data at higher Q^2 range, one needs to use perturbative

QCD and treat these ratios (at $Q^2 = \mu^2$) as inputs, i.e. boundary conditions, in the QCD evolution equations. If the numerator and denominator of the ratio have the same or almost the same factor of Q^2 dependence, then this ratio will not sensitive to the change of Q^2 .

3.3. Ellis-Jaffe sum rule

In the framework of SU(4) parton model, the first moment of the spin structure function $g_1^p(x, Q^2)$ in the proton is

$$\begin{aligned} \Gamma_1^p &\equiv \int_0^1 g_1^p(x, Q^2) dx \\ &= (4\Delta u + \Delta d + \Delta s + 4\Delta c)/18, \end{aligned} \quad (40)$$

which can be rewritten as

$$\begin{aligned} \Gamma_1^p &\equiv \int_0^1 g_1^p(x, Q^2) dx \\ &= (3a_3 + a_8 - a_{15} + 5a_0)/36 \end{aligned} \quad (41)$$

where the notations

$$a_3 = \Delta u - \Delta d, \quad a_8 = \Delta u + \Delta d - 2\Delta s \quad (42)$$

$$a_{15} = \Delta u + \Delta d + \Delta s - 3\Delta c \quad (43)$$

$$a_0 = \Delta u + \Delta d + \Delta s + \Delta c \quad (44)$$

have been introduced. Using exchange $\Delta u \leftrightarrow \Delta d$, or $a_3 \leftrightarrow -a_3$ in (41), we can obtain Γ_1^n for the neutron.

4. Numerical results and discussion.

Since the probability of charm-related splitting (9) is much smaller than u, d, s -related splittings (8), we expect that the values of parameters a and ϵ in SU(4) should be very close to those used in the SU(3) version. We choose $a = 0.143$, $\epsilon = 0.454$ [note that $a = 0.145$, $\epsilon = 0.460$ in SU(3)], and leave ϵ_c as a *variable*, then express the quark flavor and helicity contents as functions of ϵ_c . To determine the range of ϵ_c , we plot a_3 as the function of ϵ_c in Fig.1. Using the most precise data from the neutron β -decay [33], $a_3 = (G_A/G_V)_{n \rightarrow p} = 1.2670 \pm 0.0035$, we obtain $\epsilon_c \simeq 0.06 \pm 0.02$. To include other possible theoretical uncertainties arise from the model approximations, however, we prefer to introduce a larger uncertainty and take

$$\epsilon_c \simeq 0.06 \pm 0.04. \quad (45)$$

In addition to the three-parameter set $\{a, \epsilon, \epsilon_c\}$ [SU^(a)(4)], we also consider the four-parameter set: $\{a, \epsilon, \zeta', \epsilon_c\}$ [SU^(b)(4)] to show the η'^0 effect. The parameter sets used in this paper and in previous SU(3) version are listed in Table II.

Using the parameter sets SU^(a)(4) and SU^(b)(4), the flavor and spin observables in the proton are calculated and listed in Tables III and IV respectively. For comparison, we also list the existing data, the results from previous SU(3) description [27, 28], the naive quark model

(NQM) prediction, and results given by other models or analyses. Including the QCD radiative corrections, the first moments of the spin structure functions g_1^p and g_1^n are also shown in Table IV. We also calculated two reduced matrix elements F and D , and the ratios G_A/G_V of the hyperon beta decays. The results are listed in Table V. One can see that both $SU^{(a)}(4)$ and $SU^{(b)}(4)$ versions satisfactorily describe almost all the existing data. Furthermore, they give many new predictions on charm-related observables, which are in the bold type shown in Tables III and IV (for simplicity, we use $\epsilon_c = 0.06$ in $SU^{(b)}(4)$ without the uncertainty). A few main predictions selected from $SU^{(a)}(4)$ results are listed in the following:

$$2\bar{c}/\sum(q + \bar{q}) \simeq 0.011 \pm 0.008 \quad (46)$$

measures the size of the intrinsic charm in the proton,

$$\Delta c \simeq -0.009 \pm 0.006 \quad (47)$$

is the charm helicity, and

$$\Delta c/\Delta \Sigma \simeq -(0.021 \pm 0.014) \quad (48)$$

is the fraction of total quark helicity carried by the charmed quark. Several remarks are in order:

- Comparing the $SU^{(a)}(4)$ with $SU^{(b)}(4)$ predictions, one can see that there are 10-20% differences between them only for the light-flavor contents, e.g. $\bar{d}-\bar{u}$ (0.111 $\rightarrow$ 0.141), $\bar{d}/\bar{u}$ (0.71 $\rightarrow$ 0.64), $2\bar{s}/(\bar{u}+\bar{d})$ (0.66 $\rightarrow$ 0.75), and $2\bar{s}/(u+d)$ (0.118 $\rightarrow$ 0.133) [we note that both predictions are consistent with the existing data within errors], and all other predictions are almost no changes. It shows that the η^0 has minor or no effect on the spin and charm-related observables in the model. This is because first the η^0 does not have $c, \bar{c}$ components and second the $q_\uparrow$ and $q_\downarrow$ (or $\bar{q}_\uparrow$ and $\bar{q}_\downarrow$), $q = u, d, s$, appeared with equal probability in η^0 . It should be noted that the approximation $\zeta'^2 \simeq 0$ does not mean the probability of fluctuation $q \rightarrow q' + \eta^0$ ($\zeta'^2 a$) is smaller than that of $q \rightarrow q' + \bar{D}^0$ ($\epsilon_c a$). Actually, we have $\epsilon_c < \zeta'^2$. Hence we use the approximation $\zeta'^2 = 0$ in $SU^{(a)}(4)$ version only for the practical reason explained here.
- The probability a of the fluctuation $u \rightarrow d + \pi^+$ has been estimated in chiral field theory (see e.g. [22, 27])

$$a_{u \rightarrow d + \pi^+} = \int_0^1 dz \Theta(\Lambda^2 - \tau(z)) P_{u \rightarrow d + \pi^+}(z), \quad (49)$$

where

$$P_{u \rightarrow d + \pi^+}(z) = \frac{g_A^2 (m_u + m_d)^2}{32\pi^2 f_\pi^2} z \cdot \int_{-\Lambda^2}^{-\tau(z)} dt \frac{t - (m_u + m_d)^2}{(t - m_\pi^2)^2}, \quad (50)$$

and $-\tau(z) = m_u^2 z - m_d^2 z/(1-z)$. In Eq.(50), $g_A \simeq 0.75$ is the dimensionless axial-vector coupling, $f_\pi \simeq 0.093$ GeV the pion decay constant, m_u (m_d) the constituent mass of the u (d) quark, Λ the ultraviolet cutoff, and m_π is the pion mass. For $\Lambda \simeq 2.4$ GeV, one obtains $a \simeq 0.142$. For the strange quark fluctuation, $u \rightarrow s + K^+$, similar calculation with some approximation leads to the probability $\epsilon a \simeq 0.062$, which gives $\epsilon \simeq 0.44$. We assume the same formula can be used for the charm fluctuation, $u \rightarrow c + \bar{D}^0$, and obtain $\epsilon_c a \simeq 0.004$, which gives $\epsilon_c \simeq 0.03$, which is consistent with $\epsilon_c \simeq 0.06 \pm 0.04$ given in Eq.(45). Physically, the probability of splitting to the heavier mesons should be less than of splitting to the lighter ones. Hence the above estimation is quite reasonable.

- The theoretical uncertainties shown in (46)-(48) and in Tables III, IV and V arise only from the uncertainty of ϵ_c in (45). If the observable does not depend on ϵ_c , such as $\bar{d} - \bar{u}$, $\bar{d}/\bar{u}$, $2\bar{s}/(\bar{u} + \bar{d})$, etc. (these quantities depend only on $\tilde{A}$, $\tilde{B}$, and $\tilde{C}$, and not on $\tilde{D}$, i.e. independent of ϵ_c), there is no uncertainty for them. This has been shown in Table III. Two special quantities $\Delta c/c$ and $\Delta s/s$ are also independent of ϵ_c as shown in Table IV. We put a star (*) mark on some ‘data’ in Tables III and IV to denote they are model predictions or from theoretical analyses.
- The $SU(4)$ quark-meson model predicts an intrinsic charm component of the nucleon, $2\bar{c}/\sum(q + \bar{q}) \simeq 1\%$, which agrees with the predictions given in [2, 4, 12] and is also close to the those given in [3, 7] and [13, 17]. We note that the IC component is almost one order of magnitude smaller than the intrinsic strange component $2\bar{s}/\sum(q + \bar{q})$.
- Using the approach given in a previous work (see Eq. (3.6) in [27]), we can show that

$$2 \int_0^1 dx x \bar{c}(x) / \sum \int_0^1 dx x [q(x) + \bar{q}(x)] < 2\bar{c}/\sum(q + \bar{q}), \quad (51)$$

where, as defined in this paper, $\bar{c} \equiv \int_0^1 dx \bar{c}(x)$ and $\sum(q + \bar{q}) \equiv \sum \int_0^1 dx [q(x) + \bar{q}(x)]$. The l.h.s. of Eq. (51) is the fraction of the total quark momentum carried by the charm and anticharm quarks. The prediction (46) implies that this fraction is less than 1%. Assuming the quark and antiquark share about one half of the nucleon momentum, then the charm and anticharm carry about 0.5% of the nucleon momentum or less.

- From Table III, we have

$$\begin{aligned} u + \bar{u} &: d + \bar{d} : s + \bar{s} : c + \bar{c} \\ &\simeq 0.53 : 0.37 : 0.09 : 0.01. \end{aligned} \quad (52)$$

If we assume the quarks carry about 55% of the nucleon momentum, Eq. (52) implies that the fractions of the nucleon momentum carried by u -, d -, s -, and c -flavors are approximately 29.2%, 20.3%, 4.9%, and 0.6% respectively. They may compare with 31.4%, 17.8%, 4.3%, and 1.2% given by the DIS data at $Q^2=20 \text{ GeV}^2$, where the gluons carry about 45% of the nucleon momentum.

- The prediction of intrinsic charm polarization, $\Delta c \simeq -0.009 \pm 0.006$ is close to the result $\Delta c = -0.020 \pm 0.005$ given in the instanton model [6]. This might be related to the strong suppression of non-spinflip contribution in both models. Our Δc in (47) is smaller in magnitude than that given in [5, 14] ($\simeq -0.3$). However, the size of Δc given in [7] ($\simeq -5 \cdot 10^{-4}$) is even smaller. Hence further investigation in this quantity is needed.
- The ratio $\Delta c/\Delta\Sigma$ as the function of ϵ_c is plotted in Fig.2. Taking $\epsilon_c \simeq 0.06$, one has $\Delta c/\Delta\Sigma \simeq -0.021$. This is consistent with the prediction given in [6], but smaller than that given in [5]. Combining with the fractions of the light quark helicities, we have

$$\Delta u/\Delta\Sigma \simeq 2.171, \quad \Delta d/\Delta\Sigma \simeq -0.988, \quad (53)$$

$$\Delta s/\Delta\Sigma \simeq -0.162, \quad \Delta c/\Delta\Sigma \simeq -0.021,$$

one can see that the u -quark helicity is *positive* (*parallel* to the nucleon spin) and about two times larger than the total quark helicity $\Delta\Sigma$. The d -, s -, and c -helicities, however, are all *negative* (*antiparallel* to the nucleon spin), and their sizes are decrease as

$$|\Delta d| : |\Delta s| : |\Delta c| \simeq 1 : 10^{-1} : 10^{-2}. \quad (54)$$

Compare to the intrinsic strange helicity Δs , the intrinsic charm helicity is one order of magnitude smaller.

Since the hyperon β -decay data are measured at low Q^2 and our model predictions are defined at the scale $(0.2 \text{ GeV})^2 < \mu^2 < (1.0 \text{ GeV})^2$, hence we may compare them with less ambiguity. However, many data listed in Tables III and IV are coming from the DIS measurements at higher Q^2 range. To make a meaningful comparison of model predictions with these data, we have to discuss the

Q^2 dependence of these observables. For spin observable, as we mentioned in section III of [27] that the model predictions, e.g. Δu , Δd , Δs , etc. are compared with the (factorization) scheme-independent DIS observables $a_q(Q^2) \equiv \Delta q - [\alpha_s(Q^2)/2\pi]\Delta G(Q^2)$ ($q = u, d, s$) at the same Q^2 scale [i.e. $(0.2 \text{ GeV})^2 < \mu^2 < (1.0 \text{ GeV})^2$], where $\Delta G(Q^2)$ is the helicity of the gluon, and the axial charge $a_q(Q^2)$ is defined in the Adler-Bardeen scheme. Although $a_q(Q^2)$ is independent of Q^2 at the leading order and changes very slowly with Q^2 at NLO, we still need to assume the perturbative QCD can be used down to the scale μ^2 . The perturbative QCD evolution approach has been successfully used down to $Q^2 \simeq 0.23 \text{ GeV}^2$ [34], it is not clear, however, if the approach still hold below this Q^2 .

Finally, in the quark meson fluctuation model, it is possible to include the contributions come from quark splittings to the vector mesons such as K^* , ρ , etc. [1^- nonet in SU(3) and 1^- 15-plet in SU(4)]. However, it will change the formalism completely and is far beyond the goal of this paper. For simplicity and consistency, we only discuss the contributions of 0^- pseudoscalar mesons at this moment and defer the discussion of possible vector meson fluctuations to a later time.

In summary, we have calculated the intrinsic charm contribution in the SU(4) quark meson model with symmetry breaking. Despite the approximations and possible theoretical uncertainties, the overall agreement between the predictions and the existing data seems to be quite satisfactory considering the model is *simple* and has only a *few* parameters. The model also leads to many *new* predictions on observables explicitly related to the charm or anticharm. These observables are *zero*, e.g. $2\bar{c}/\sum(q+\bar{q})$, Δc , etc. or *indefinite*, e.g. $\Delta c/c$ and $c_\uparrow/c_\downarrow$, in the SU(3) description. We hope that these predictions can be tested by the analyses of the DIS data on polarized and unpolarized charm productions in the near future.

Acknowledgments



This work was supported in part by the U.S. DOE Grant, the Institute of Nuclear and Particle Physics, Department of Physics, University of Virginia, and the Commonwealth of Virginia.

[1] S. J. Brodsky, P. Hoyer, C. Peterson and N. Sakai, Phys. Lett. **B93**, 451 (1980); S. J. Brodsky, C. Peterson, and N. Sakai, Phys. Rev. **D23**, 2745 (1980).
[2] J. F. Donoghue and E. Golowich, Phys. Rev. **D15**, 3421 (1977); H. He, X. Zhang, and Y. Zhao, High Energy Phys. Nucl. Phys. **7**, 626 (1983).
[3] T. Hatsuda, and T. Kunihiro, Phys. Rep. **247**, 221

(1994).
[4] F. S. Navarra, M. Nielson, C. A. A. Nunes, and M. Teixeira, Phys. Rev. **D54**, 842 (1996).
[5] A. Blotz and E. Shuryak, Phys. Lett. **B439**, 415 (1998).
[6] F. Araki, M. Musakhanov and H. Toki, hep-ph/9808290.
[7] M. V. Polyakov, A. Schafer, and O. V. Teryaev, Phys. Rev. **D60**, 051502 (1999); M. Franz, M. V. Polyakov,

- [0] and K. Goeke, Phys. Rev. **D62**, 074024 (2000).
- [8] E. Hoffmann and R. Moore, Z. Phys. **C20**, 71 (1983).
- [9] B. W. Harris, J. Smith and R. Vogt, Nucl. Phys. **B461**, 181 (1996).
- [10] R. Vogt and S. J. Brodsky, Nucl. Phys. **B478**, 311 (1996).
- [11] G. Ingelman and M. Thunman, Z. Phys. **C73**, 505 (1997).
- [12] Y. A. Golubkov, *Phys. Atom. Nucl.* **63**, 606 (2000); hep-ph/9811218.
- [13] S. J. Brodsky, and I. Schmidt, Phys. Rev. **D43**, 179 (1991).
- [14] I. Halperin and A. Zhitnitsky, Phys. Rev. **D56**, 7247 (1997).
- [15] J. F. Gunion and R. Vogt, hep-ph/9706252.
- [16] W. Melnitchouk and A. T. Thomas, Phys. Lett. **B414**, 134 (1997).
- [17] Steffens, W. Melnitchouk and A. T. Thomas, Eur. Phys. J. **C11**, 673 (1999).
- [18] J. Blumlein and W. L. van Neerven, Phys. Lett. **B450**, 417 (1999).
- [19] H1 Collaboration, C. Adloff *et al.* Z. Phys. **C72**, 593 (1996).
- [20] ZEUS Collaboration, J. Breitweg *et al.* Phys. Lett. **B407**, 402 (1997).
- [21] R. Vogt, Prog. Part. Nucl. Phys. **45**, S105 (2000).
- [22] E. J. Eichten, I. Hinchliffe and C. Quigg, Phys. Rev. **D45**, 2269 (1992).
- [23] T. P. Cheng and L.-F. Li, Phys. Rev. Lett. **74**, 2872 (1995).
- [24] X. Song, J. S. McCarthy and H. J. Weber, Phys. Rev. **D55**, 2624 (1997).
- [25] H. J. Weber, X. Song, and M. Kirchbach, Mod. Phys. Lett. **A12**, 729 (1997).
- [26] T. P. Cheng and L.-F. Li, Phys. Rev. **D57**, 344 (1998).
- [27] X. Song, Phys. Rev. **D57**, 4114 (1998).
- [28] X. Song, Int. J. Mod. Phys. **A16**, 2673 (2001).
- [29] X. Song, Internal Report, ICTP, June (1998).
- [30] S. L. Glashow *et al.*, Phys. Rev. **D2**, 2185 (1970).
- [31] Presumably one should also consider the contributions coming from other possible nonperturbative mechanisms, such as $gg \rightarrow q\bar{q}$, which are not included in this paper. However, it is not clear whether these nonperturbative mechanisms are independent or not.
- [32] K. Abe, *et al.*, Phys. Rev. Lett. **79**, 26 (1997).
- [33] Particle Data Group, D. E. Groom, *et al.*, Eur. Phys. J. **C15**, 1 (2000).
- [34] M. Gluck, E. Reya, and A. Vogt, Z. Phys. **C47**, 433 (1995).
- [35] J. C. Peng, *et al.*, E866/NuSea Collaboration, Phys. Rev. **D58**, 092004 (1998).
- [36] P. Amaudruz *et al.*, Phys. Rev. Lett. **66**, 2712 (1991); M. Arneodo *et al.*, Phys. Rev. **D50**, R1, (1994).
- [37] A. Baldit, *et al.*, NA51 Collaboration, Phys. Lett. **B332**, 244 (1994).
- [38] A. O. Bazarko, *et al.*, Z. Phys. **C65**, 189 (1995).
- [39] J. Gasser, H. Leutwyler and M. E. Saino, Phys. Lett. **B253**, 252 (1991).
- [40] S. J. Dong, J. F. Lagae and K. F. Liu, Phys. Rev. **D54**, 5496 (1996).
- [41] P. Adams, *et al.*, Phys. Rev. **D56**, 5330 (1997).
- [42] B. Adeva, *et al.*, Phys. Lett. **B369**, 93 (1996).

Figures

FIG. 1: a_3 as the function of ϵ_c .FIG. 2: Intrinsic charm quark polarization in the proton as the function of ϵ_c .

Tables

TABLE I: The probabilities $P_{q_{\uparrow}}(q'_{\uparrow,\downarrow}, \bar{q}'_{\uparrow,\downarrow})$ and $P_{q_{\downarrow}}(q'_{\uparrow,\downarrow}, \bar{q}'_{\uparrow,\downarrow})$.

q'	$P_{u_{\uparrow}}(q'_{\uparrow,\downarrow})$	$P_{d_{\uparrow}}(q'_{\uparrow,\downarrow})$	$P_{s_{\uparrow}}(q'_{\uparrow,\downarrow})$
$u_{\uparrow}$	$1 - (1 + \epsilon + \epsilon_c + 2f)a/2 + (1 - \tilde{A})^2 a/2$	$\tilde{A}^2 a/2$	$\tilde{B}^2 a/2$
$u_{\downarrow}$	$(1 + \epsilon + \epsilon_c + 2f)a/2 + (1 - \tilde{A})^2 a/2$	$a + \tilde{A}^2 a/2$	$\epsilon a + \tilde{B}^2 a/2$
$d_{\uparrow}$	$\tilde{A}^2 a/2$	$1 - (1 + \epsilon + \epsilon_c + 2f)a/2 + (1 - \tilde{A})^2 a/2$	$\tilde{B}^2 a/2$
$d_{\downarrow}$	$a + \tilde{A}^2 a/2$	$(1 + \epsilon + \epsilon_c + 2f)a/2 + (1 - \tilde{A})^2 a/2$	$\epsilon a + \tilde{B}^2 a/2$
$s_{\uparrow}$	$\tilde{B}^2 a/2$	$\tilde{B}^2 a/2$	$1 - (\epsilon + f_s + \epsilon_c/2)a + \tilde{C}^2 a/2$
$s_{\downarrow}$	$\epsilon a + \tilde{B}^2 a/2$	$\epsilon a + \tilde{B}^2 a/2$	$(\epsilon + f_s + \epsilon_c/2)a + \tilde{C}^2 a/2$
$c_{\uparrow}$	$\tilde{D}^2 a/2$	$\tilde{D}^2 a/2$	$\tilde{D}^2 a/2$
$c_{\downarrow}$	$\epsilon_c a + \tilde{D}^2 a/2$	$\epsilon_c a + \tilde{D}^2 a/2$	$\epsilon_c a + \tilde{D}^2 a/2$
$\bar{u}_{\uparrow,\downarrow}$	$(1 - \tilde{A})^2 a/2$	$(1 + \tilde{A}^2)a/2$	$(\epsilon + \tilde{B}^2)a/2$
$\bar{d}_{\uparrow,\downarrow}$	$(1 + \tilde{A}^2)a/2$	$(1 - \tilde{A})^2 a/2$	$(\epsilon + \tilde{B}^2)a/2$
$\bar{s}_{\uparrow,\downarrow}$	$(\epsilon + \tilde{B}^2)a/2$	$(\epsilon + \tilde{B}^2)a/2$	$\tilde{C}^2 a/18$
$\bar{c}_{\uparrow,\downarrow}$	$(\epsilon_c + \tilde{D}^2)a/2$	$(\epsilon_c + \tilde{D}^2)a/2$	$(\epsilon_c + \tilde{D}^2)a/2$

TABLE II: Parameter sets for different symmetry-breaking schemes

Model	a	ϵ	$\zeta'^2_{SU(3)}$	ϵ_c
SU ^(a) (4)	0.143	0.454	—	0.06 ± 0.04
SU ^(b) (4)	0.143	0.454	0.1	0.06
SU(3)[28]	0.145	0.460	0.1	—

TABLE III: Quark Flavor Observables.

Quantity	Data	SU ^(a) (4)	SU ^(b) (4)	SU(3)[28]	NQM
$\bar{d} - \bar{u}$	$0.110 \pm 0.018[35]$ $0.147 \pm 0.039[36]$	0.111	0.141	0.143	0
$\bar{u}/\bar{d}$	$[\bar{u}(x)/\bar{d}(x)]_{0.1 < x < 0.2} = 0.67 \pm 0.06[35]$ $[\bar{u}(x)/\bar{d}(x)]_{x=0.18} = 0.51 \pm 0.06[37]$	0.71	0.64	0.64	—
$2\bar{s}/(\bar{u} + \bar{d})$	$< 2x\bar{s}(x) > / < x(\bar{u}(x) + \bar{d}(x)) >$ $= 0.477 \pm 0.051[38]$	0.66	0.75	0.76	—
$2\bar{c}/(\bar{u} + \bar{d})$	—	0.083 ± 0.055	0.085	0	—
$2\bar{s}/(u + d)$	$< 2x\bar{s}(x) > / < x(u(x) + d(x)) >$ $= 0.099 \pm 0.009[38]$	0.118	0.133	0.136	0
$2\bar{c}/(u + d)$	—	0.015 ± 0.010	0.015	0	0
$(u + \bar{u})/\sum(q + \bar{q})$	—	0.530 ± 0.004	0.519	0.523	2/3
$(d + \bar{d})/\sum(q + \bar{q})$	—	0.368 ± 0.003	0.370	0.374	1/3
$(s + \bar{s})/\sum(q + \bar{q})$	$< 2x\bar{s}(x) > / \sum < x(q(x) + \bar{q}(x)) >$ $= 0.076 \pm 0.022[38]$ $= 0.10 \pm 0.06[39]$ $= 0.15 \pm 0.03[40]$	0.090 ± 0.001	0.100	0.103	0
$(c + \bar{c})/\sum(q + \bar{q})$	0.03 [13]* 0.02 [2]* 0.01 [12]* 0.009 [4]* 0.005 [3, 7]* ≤ 0.004 [17]*	0.011 ± 0.008	0.011	0	0
$\sum \bar{q}/\sum q$	$\sum < x\bar{q}(x) > / \sum < xq(x) >$ $= 0.245 \pm 0.005[38]$	0.230 ± 0.004	0.233	0.231	0

TABLE IV: Quark Spin Observables

Quantity	Data	SU ^(a) (4)	SU ^(b) (4)	SU(3)[28]	NQM
Δu	$0.85 \pm 0.04[41]$	0.871 ± 0.009	0.859	0.863	4/3
Δd	$-0.41 \pm 0.04[41]$	-0.397 ± 0.002	-0.393	-0.397	-1/3
Δs	$-0.07 \pm 0.04[41]$	-0.065 ± 0.000	-0.065	-0.067	0
Δc	-0.020 ± 0.004 [6]* -0.3 [5]* $-5 \cdot 10^{-4}$ [7]*	-0.009 ± 0.006	-0.009	0	0
$\Delta\Sigma/2$	$0.19 \pm 0.06[41]$	0.200 ± 0.006	0.196	0.200	1/2
$\Delta\bar{u}, \Delta\bar{d}$	$-0.02 \pm 0.11[42]$	0	0	0	0
$\Delta\bar{s}, \Delta\bar{c}$	—	0	0	0	0
$\Delta u/\Delta\Sigma$	—	2.171 ± 0.043	2.192	2.162	4/3
$\Delta d/\Delta\Sigma$	—	-0.988 ± 0.024	-1.004	-0.994	-1/3
$\Delta s/\Delta\Sigma$	—	-0.162 ± 0.005	-0.166	-0.167	0
$\Delta c/\Delta\Sigma$	-0.08 ± 0.01 [5]* -0.033 [6]*	-0.021 ± 0.014	-0.022	0	0
$\Delta u/u$	—	0.383 ± 0.003	0.381	0.383	2/3
$\Delta d/d$	—	-0.287 ± 0.001	-0.283	-0.284	-1/3
$\Delta s/s$	—	-3/10	-0.269	-0.269	—
$\Delta c/c$	—	-16/51	-16/51	—	—
$u_{\uparrow}/u_{\downarrow}$	—	2.241 ± 0.012	2.231	2.241	5
$d_{\uparrow}/d_{\downarrow}$	—	0.554 ± 0.001	0.559	0.558	1/2
$s_{\uparrow}/s_{\downarrow}$	—	7/13	0.576	0.576	—
$c_{\uparrow}/c_{\downarrow}$	—	35/67	35/67	—	—

Γ_1^p	$0.136 \pm 0.016[41]$	0.143 ± 0.002	0.141	0.142	5/18
Γ_1^n	$-0.041 \pm 0.007[32]$	-0.042 ± 0.001	-0.043	-0.042	0
a_3	$1.2670 \pm 0.0035[33]$	1.268 ± 0.010	1.252	1.260	5/3
a_8	$0.579 \pm 0.025[33]$	0.605 ± 0.006	0.595	0.600	1

TABLE V: F, D and G_A/G_V ratios of the hyperon beta decays. [a]: This is the best fit to four measured (G_A/G_V) ($\chi^2 \simeq 1.96$) under the constraint F+D=1.267.

Quantity	Data	SU ^(a) (4)	SU ^(b) (4)	SU(3)[28]	NQM
F+D	1.267[a]	1.268 ± 0.010	1.252	1.260	5/3
F	0.463[a]	0.468 ± 0.004	0.462	0.465	2/3
D	0.804[a]	0.800 ± 0.006	0.790	0.796	1
F/D	0.576[a]	0.585 ± 0.007	0.585	0.585	2/3
$(G_A/G_V)_{n \rightarrow p}$	$1.2670 \pm 0.0035[33]$	1.268 ± 0.010	1.252	1.260	5/3
$(G_A/G_V)_{\Lambda \rightarrow p}$	$0.718 \pm 0.015[33]$	0.735 ± 0.007	0.725	0.730	1
$(G_A/G_V)_{\Sigma^- \rightarrow n}$	$-0.340 \pm 0.017[33]$	-0.332 ± 0.002	-0.328	-0.330	-1/3
$(G_A/G_V)_{\Xi^- \rightarrow \Lambda}$	$0.25 \pm 0.05[33]$	0.202 ± 0.002	0.198	0.200	1/3